

(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 078 923 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
08.03.2006 Bulletin 2006/10

(51) Int Cl.:
C07D 333/54 (2006.01) C07D 417/14 (2006.01)

(21) Application number: **00115890.6**

(22) Date of filing: **25.07.2000**

(54) Process for the preparation of benzothiophene derivatives

Verfahren zur Herstellung von Benzothiophen-Derivaten

Procédé de préparation de dérivés de benzothiophène

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(30) Priority: **02.08.1999 EP 99115222**

(43) Date of publication of application:
28.02.2001 Bulletin 2001/09

(73) Proprietor: **F. HOFFMANN-LA ROCHE AG
4070 Basel (CH)**

(72) Inventors:
• **Junghans, Bernd
68535 Edingen-Neckarhausen (DE)**
• **Scalone, Michelangelo
4127 Birsfelden (CH)**
• **Zeibig, Thomas Albert
67112 Mutterstadt (DE)**

(74) Representative: **Witte, Hubert et al
Grenzacherstrasse, 124
4070 Basel (CH)**

(56) References cited:
WO-A-94/27995

- **M IWASAKI ET AL: "Palladium-Catalyzed
Cyclocarbonylation of 3-(Heteroaryl)allyl
Acetates" JOURNAL OF ORGANIC CHEMISTRY,
US,AMERICAN CHEMICAL SOCIETY. EASTON,
vol. 56, no. 5, 1991, pages 1922-1927,
XP002082809 ISSN: 0022-3263**

Remarks:

The file contains technical information submitted after
the application was filed and not included in this
specification

EP 1 078 923 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).


Description

[0001] The present invention is concerned with a novel process for the preparation of benzothiophene derivatives, especially with the preparation of 4-hydroxybenzothiophene. 4-Hydroxybenzothiophene is a building block for pharmaceutically active compounds, e.g. 5-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]-7-benzothiophenylmethyl]-2,4-thiazolidinedione. This compound is known in the art and is described for example in International Patent Application WO 94/27995. It is especially useful for prophylaxis and treatment of diabetes mellitus type I and II.

[0002] Methods for the preparation of 4-hydroxybenzothiophene have been described by Iwasaki et al. (1991) J. Org. Chem. 1991. 56, 1922. Here a cyclocarbonylation of a primary allylacetate is performed in presence of a high catalyst loading. Further, this process is characterized by at least five process steps which in part require extreme reaction conditions.


[0003] Surprisingly it has been found that using the process according to the present invention 4-hydroxybenzothiophene can be prepared with less process steps under moderate conditions with an outstanding yield.

[0004] The process according to the present invention refers to the 4-hydroxybenzothiophene of Formula I



I

comprising cyclocarbonylation of a compound of Formula II



II

wherein

-OR is a group of formulae -O-(CO)-R', -O-(CO)-O-R'', or -O-(PO)-(OR'')₂, wherein R' is alkyl, perfluoro-C₁₋₂₀-alkyl, aryl, R'' is alkyl, aryl or benzyl or the group -OR is halogen or an aryloxy group; followed by saponification.

[0005] This process provides an efficient cyclocarbonylation reaction under mild conditions. In addition, substrates for the cyclocarbonylation reaction (compound of Formula II) do not need to be purified, e.g. by distillation, but can be used as "crude" material.

[0006] According to the present invention, the term "cyclocarbonylation" refers to an introduction of a carbonyl group coupled to the formation of a cyclic ring structure.

[0007] The term "saponification" refers to the hydrolysis of an ester under acid or basic, preferably basic conditions.

[0008] The term "transition metal compound" refers to a metal-phosphine complex compound wherein the term metal refers to Pd, Pt, Ru, Co; Rh or Ni, preferably Pd.

[0009] The term "ligand" refers to phosphine, arsine or stibine derivatives, preferable phosphine derivatives, of general formulae P(R¹)(R²)(R³), (R¹)(R²)P-(X)-P(R¹)(R²), As(R¹)(R²)(R³) or Sb(R¹)(R²)(R³), preferably P(R¹)(R²)(R³), wherein R¹, R², and R³ are defined below.

[0010] The term "alkyl" refers to a branched or straight chain monovalent alkyl radical of one to nine carbon atoms (unless otherwise indicated), preferably one to four (lower) carbon atoms. This term is further exemplified by such radicals as methyl, ethyl, n-propyl, isopropyl, i-butyl, n-butyl, t-butyl and the like.

[0011] The term "aryl" refers to a monovalent carbocyclic aromatic radical, e.g. phenyl, optionally substituted, independently, with halogen, lower-alkyl, lower-alkoxy, lower-alkylenedioxy, carboxy, trifluoromethyl and the like.


[0012] The term "aryloxy", signifies a group of the formula aryl-O- in which the term "aryl" has the significance given above. Phenylloxy is an example of such an aryloxy group.

[0013] The term "alkoxy", alone or in combination, signifies a group of the formula alkyl-O- in which the term "alkyl" has the significance given above, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec.butoxy and tert.butoxy, preferably methoxy and ethoxy.

[0014] The term "alkylenedioxy" refers to C₁₋₃-alkyl-dioxy groups, such as methylenedioxy, ethylenedioxy or propylenedioxy.


[0015] The term "halogen" refers to fluorine, chlorine, and bromine.

[0016] In more detail, the present invention refers to a process for the preparation of compounds of Formula I



I

comprising cyclocarbonylation of a compound of Formula II



II

wherein

-OR is a group of formulae -O-(CO)-R', -O-(CO)-O-R'' or -O-(PO)-(OR'')₂, wherein R' is alkyl, perfluoro-C₁₋₂₀-alkyl, aryl, R'' is alkyl, aryl or benzyl or the group -OR is halogen or an aryloxy group; followed by saponification.

[0017] In a preferred embodiment of the invention, the cyclocarbonylation reaction is carried out in the presence of a base and a catalyst comprising a transition metal compound and a ligand.


[0018] Transition metal compounds useful for the process of the present invention comprise salts of Pd, Pt, Ru, Co, Rh- or Ni and also includes Pd/C. The use of transition metal compounds as catalysts has been described for example in Matsuzaka et al. (1988) J. Org. Chem. 53, 3832. Preferred transition metal compounds are salts of palladium, e.g. Pd(OAc)₂, Pd₂dba₃, PdCl₂, Pd₂Cl₂(π-allyl)₂, PdCl₂(NCMe)₂, [Pd(NCMe)₄](BF₄)₂, and most preferably Pd(OAc)₂. The mentioned catalysts are known in the art (e.g. US Patent No. 5,380,861; "Carbonylation, Direct Synthesis of Carbonyl Compounds", H.M. Colquhoun, D.J. Thompson, M.V. Trigg, Plenum Press, 1991) and/or are commercially available (e.g. from Fluka, Buchs, Switzerland or Strem Chemicals, Kehl, Germany).

[0019] The ligand of the transition metal compound in the catalyst may be selected from a group consisting of phosphine, arsine or stibine derivatives, preferable phosphine derivatives of general formulae P(R¹)(R²)(R³), (R¹)(R²)P-(X)-P(R¹)(R²), As(R¹)(R²)(R³) or Sb(R¹)(R²)(R³), preferably P(R¹)(R²)(R³), wherein X, R¹, R², and R³ are defined below.

[0020] Especially suitable ligands are chiral and non-chiral mono- and diphosphorus compounds for example described in Houben-Weyl, "Methoden der organischen Chemie", vol. E1, page 106 et seq. Georg Thieme Verlag Stuttgart, 1982, and Aspects Homog. Catal., 4, 145-202 (1981), especially those of the formulae



wherein R¹, R² and R³ each independently are C₁₋₈-alkyl, cyclohexyl, benzyl, naphthyl, 2- or 3-pyrrolyl, 2- or 3-furyl, 2- or 3-thiophenyl, 2- or 3- or 4-pyridyl, phenyl or phenyl which is substituted by C₁₋₄-alkyl, C₁₋₄-alkoxy, halogen, trifluoromethyl, lower alkyldenedioxy or phenyl and X is binaphthyl, 6,6'-dimethyl- or 6,6'-dimethoxybiphenyl-2,2'-diyl, or one of the groups -(CH₂)_n-, -CH₂CH₂-P(C₆H₅)-CH₂CH₂-,



IV

and n is a number of 1 - 8.

[0021] Examples of suitable phosphorus ligands are shown in Scheme 1.

5

10

15

20

25

30







35

40


45

50

55

Scheme 1:

[0022] The most preferred phosphorus ligands are triphenylphosphine,



[0023] The preparation of a transition metal complex is explained in more detail for the corresponding palladium-phosphine complex: The palladium-phosphine complex compound is conveniently formed in situ from a palladium component and a phosphine ligand. These palladium components is for example metallic palladium, which is optionally supported on a carrier material such as carbon, or a complex or a salt of 0-, 2- or 4-valent palladium such as palladium-bis (dibenzylideneacetone), palladium chloride, palladium acetate and the like. For the in situ preparation, the phosphorus ligand/transition metal compound ratio (mol/mol; P/Pd) amounts to about 0.1 : 1 to 100 : 1, preferably to about 6 : 1 to 15 : 1. Suitable phosphine ligands are for example chiral and non-chiral mono- and diphosphorus compounds such as are described in Houben-Weyl, Methoden der organischen Chemie, volume E1, page 106 et. seq. Georg Thieme Verlag Stuttgart, 1982, and Aspects Homog. Catal., 4, 145 - 202 (1981), especially those described above.

[0024] For the in situ preparation of the palladium-phosphine complex compound palladium-(II) chloride or palladium-(II) acetate, palladium-dichloro-bis(acetonitrile) and a bis(diphenylphosphino)alkane may be used.

[0025] Further, the process of the present invention comprises the use of bases for the cyclocarbonylation reaction like tertiary bases such as tri-alkyl-amines, di-alkyl-aryl-amines, pyridines, alkyl-N-piperidines, and for example inorganic bases such as NaOH, KOH or salts of carbonic acids. Examples are (alkyl)₃amines, e.g. triethylamine, ethyl-diisopropyl-amine, pyridine, N-methyl-piperidine, sodium hydrogen carbonate, potassium hydrogen carbonate, di-sodium carbonate, etc. The preferred base is triethylamine.

[0026] Solvents for the above reaction are known to skilled persons. Preferred solvents are aromatic solvents, e.g. toluene, xylene, benzene, halogenated hydrocarbons, e.g. CH₂Cl₂, nitriles, e.g. acetonitrile, ester, e.g. ethylacetate, amides, e.g. DMF, ether, e.g. THF, dioxane, urethanes, e.g. TMU, sulfoxides, e.g. DMSO, and mixtures thereof. The preferred solvent is toluene.

[0027] The reaction conditions for the above carbonylation reaction can vary to a certain extent.

[0028] The temperature can vary between 40°C and 170°C, preferably between 60 - 120°C, and most preferably the reaction is performed at about 90°C.

[0029] The substrate/catalyst ratio (mol/mol; S/Pd) amounts to 1 to 10000, preferably 100 to 5000, more preferably 1000 to 2000 and most preferably 1200 to 1500.


[0030] For the in situ preparation, the above mentioned phosphorus ligand/transition metal compound ratio (mol/mol; P/Pd) amounts to 0.1 : 1 to 100 : 1, preferably 6 : 1 to 15 : 1.

[0031] The upper limit for the carbon monoxide (CO) pressure is only limited by the specification of the autoclave used. For the lower pressure limit the carbonylation reaction would work even with a CO pressure of 1 bar. Preferably, the CO pressure is about 20 to 70 bar, more preferably 35 to 60 bar.

[0032] Surprisingly it has been found that the "crude" compound of Formula II can be used for the preparation of the compound of Formula I. A preparation of a crude material is performed by collecting the compound of Formula II, e.g. 1-(2-thienyl)allyl acetate, with an organic solvent and drying without further purification. The preparation of this material is exemplified in Example 1. Example 2 B shows the use of the crude starting material for the preparation of the compound of Formula I.

[0033] The cyclocarbonylation reaction is followed by saponification. Conditions for saponification reactions are known in the art and described for example in "Practical Organic Chemistry", A.I. Vogel, Longmans Ed., 1967, p. 390 - 393. In a preferred embodiment of the present invention, the saponification is carried out in a biphasic mixture of aqueous sodium hydroxide and toluene or in an homogeneous mixture of sodium methylate in methanol.

[0034] Compounds of Formula II may be prepared by methods known in the art, for example by reaction of a thiophene carbaldehyde of Formula III (illustrated in Scheme 2 a; commercially available, Fluka, Aldrich).




with a vinyl-metal-X reagent, with -metal-X being -MgCl, -MgBr, -MgI or -Li, preferably -MgCl or -MgBr, followed by reaction with an acid derivative. Other allyl compounds, e.g. the corresponding allyl halogenides or allyl trialkylammonium salts, are also suitable reagents. The acid derivative can be selected from a group consisting of compounds of formulae, (R'-CO)₂O, R''O-(CO)-Cl, Cl-(PO)(OR'')₂, R'-(CO)-Hal wherein R' is alkyl, perfluoro-C₁₋₂₀-alkyl, aryl, R'' is alkyl, aryl or benzyl and Hal is Cl or Br. The preferred acid derivative is (R'-CO)₂O, and here especially the acetic anhydride. The most preferred the vinyl-metal-X-reagents are vinylmagnesium chloride or vinylmagnesium bromide.

[0035] In the most preferred embodiment of the present invention, the compound of Formula II is prepared by reaction of vinylmagnesium chloride followed by reaction with acetic anhydride as shown in scheme 2, variant a).

[0036] Additional methods for the preparation of compound III are summarized in scheme 2.


Scheme 2:



[0037] The compound of Formula I is useful for the preparation of pharmaceutically active substances, e.g. 5-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]-7-benzothiophenylmethyl]-2,4-thiazolidinedione and its salts, especially the corresponding sodium salt. A process for the preparation of this compound has been described for example in International Patent Application WO 98/42704.

[0038] In addition, the compounds may be prepared according to the following processes:

[0039] In a first step the compound of Formula I may be converted into 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole by reaction with a mesylate of Formula V



under basic conditions. The reaction may be performed in solvents like DMF with for example sodium carbonate, potassium carbonate or cesium carbonate, preferably potassium carbonate; or in THF with KtBu; or in toluene and KOH with phase transfer catalysts.

[0040] The above process may be followed by a nitration reaction of 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole to give 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole. Normally nitric acid is used for the nitration reaction which may be performed at room temperature to about 50 °C, preferably room temperature.

[0041] The 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole obtained by the above process may be converted into 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole by hydrogenation. The conditions of the hydrogenation reaction (H_2 /Raney nickel) are known in the art. Hydrogen pressure may be 1 to 10 bar, preferably 1 bar.

[0042] The above process may be continued by the reaction of 5-methyl-4-[2-(7-aminobenzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole with $H\text{Hal}/\text{NaNO}_2$ followed by reaction with $\text{CH}=\text{CHCOOCH}_3/\text{Cu(I)Hal}$, wherein Hal is Br or Cl, preferably Br. The reaction product in case of Hal is Br is methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate.


[0043] The reaction of methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate with thiourea will produce 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methyl-thiazolidine-4-one. The reaction is normally performed in alkylalcohols like ethanol.

[0044] This compound (2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methyl-thiazolidine-4-one) may then be converted into 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione by reaction under acid conditions. The reaction may be performed at 1 - 4 bar, preferably at bar. Acidic conditions are provided by an organic or inorganic acid in an appropriate solvent, e.g. HCl/ethanol.


[0045] The reaction may be optionally continued by conversion of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione in a corresponding salt, preferably the sodium salt (5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt) by reaction under basic conditions, preferably with NaOH in THF.

[0046] A further embodiment of the invention comprises a process for the preparation of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione and/or of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione sodium salt comprising

a) conversion of a compound of Formula I into 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole by reaction of a compound of Formula I



with a mesylate of Formula V



under basic conditions; followed by

b) nitration of 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole to give 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole;

c) hydrogenation of 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole to give 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole; followed by

d) reaction of 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole with $H\text{Hal}/\text{NaNO}_2$ and $\text{CH}=\text{CHCOOCH}_3/\text{Cu(I)Hal}$, wherein Hal is Br or Cl to give methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate; followed by

e) reaction of methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate with thiourea to give 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methylthiazolidine-4-one; followed by

f) reaction of 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methyl-thiazolidine-4-one under acid conditions to give 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione;

g) and optionally followed by reaction of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione under basic conditions to give 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt.

[0047] The invention further comprises the use of any of the above described processes for the preparation of 5-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]-7-benzothiophenylmethyl]-2,4-thiazolidinedione and 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt.

[0048] A further embodiment of the present invention comprises the compound 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt.

[0049] The following examples shall illustrate preferred embodiments of the present invention.

EXAMPLES

Example 1

1-(2-Thienyl)allyl Acetate

[0050] A 1.514-necked glass flask equipped with a mechanical stirrer, a thermometer and an argon inlet was charged with 112.2 g of 2-thiophenecarbaldehyde (1.00 mol) and 100 ml of THF and to the resulting solution was added dropwise at -20°C within 1.2 h 650 ml of vinylmagnesium chloride 1.7 M solution in THF. The temperature during the addition was kept between -20 and -25°C with aid of an acetone/dry ice bath, then increased to 0°C during 35 min and kept at this temperature for 20 min. To the resulting brown suspension was added at ca. 0° within 40 min 132.7 g of acetic anhydride (1.30 mol). The cooling bath was removed and after stirring for 1 h 400 ml of deionized water was added at 10-15°C within 20 min. The biphasic yellow-brown mixture was stirred for an additional 1 h at room temperature and transferred to a separatory funnel with aid of 500 ml of hexane. The brown aqueous phase was separated and extracted with 400 ml of hexane. The combined organic phases were washed with 3 x 200 ml deionized water, dried (Na₂SO₄, 15 min stirring) and rotatory evaporated (T_{bath} 35°, 12 mbar, 1 h). Material of this quality is defined as "crude" and is also suitable for cyclocarbonylation (see Example 2 B). The orange-brown oil (199.7 g) was distilled in an apparatus consisting of a 500-ml two-necked round-bottomed flask, a distillation head with water-cooling and a fraction sampler. A forerun containing low-boiling components (yellowish oil) was collected at T_{head} between room temperature and 55°C and 0.5-0.6 mbar, the main fraction was collected at T_{head} of 59-62°C (T_{pot} 63-67°C) and 0.4 mbar.

Yield: 161.53 g (88.6%) of 1-(2-thienyl)allyl acetate as a slightly yellow oil.

Example 2 A

4-Hydroxybenzothiophene

[0051] An autoclave was charged under an argon flow with 27.34 g of 1-(2-thienyl)allyl acetate (0.150 mol, distilled), 28.4 ml of acetic anhydride (30.6 g, 0.30 mol), 42.0 ml of triethylamine (30.7 g, 0.30 mol), 23.6 mg of palladium acetate (0.105 mmol) and 0.264 g of triphenylphosphine (1.00 mmol), all with aid of 53 ml of toluene. Then the autoclave was sealed, evacuated twice under slow stirring (150 rpm) to 0.2 bar and pressurized with 8 bar of argon, then pressurized three times with 20 bar of carbon monoxide and vented, and finally pressurized with 50 bar of carbon monoxide. The reaction mixture was stirred (500 rpm) and heated at 120°C and the carbonylation carried out at 50 bar constant total pressure for 6 h. After cooling, the autoclave was vented and the CO atmosphere was exchanged by evacuating to ca. 0.2 bar and pressurizing 8 bar of argon four times. The resulting dark solution was poured into a 0.5 l flask containing 120 ml of ice water and the biphasic solution was stirred for 1 h at room temperature. The aqueous phase was extracted in a separatory funnel with 80 ml of toluene and then the combined organic phases were washed with 3 x 30 ml, with a total of 90 ml of deionized water and reduced to a total weight of 46 g by rotary evaporation (50°C/60 mbar).

[0052] The residue containing the crude acetate was transferred to a 0.351 glass flask under argon with aid of 25 ml of toluene. After addition of 82 ml of 4N sodium hydroxide (328 mmol) the mixture was stirred intensively (1200 rpm) at 50° for 1.5 h and then after cooling transferred in a 0.51 separatory funnel. After removal of the organic layer, the dark aqueous phase was extracted with 80 ml of toluene and the combined organic phases were back-extracted with 2 x 20 ml, a total of 40 ml of deionized water. The combined aqueous phases were treated with 1.0 g of charcoal, stirred at room temperature for 5 min under argon and filtered through a Speedex layer. The filter cake was rinsed three times with 20 ml, a total of 60 ml of deionized water. The clear brown, combined phases were concentrated until no more

toluene distilled, then after cooling to 5°C in an ice bath 75 ml of 25% HCl were added under argon during 35 min, whereas the temperature was kept under 15° with aid of an ice bath. The resulting thick crystalline suspension was stirred for 1 h in an ice bath (internal temperature 2-3°C) and filtered on a sintered glass filter. The filter cake was washed three times with 50 ml, a total of 150 ml of ice-cold water and dried on the rotavapor at 50°C/l mbar to constant weight.

5 Yield: 18.9 g (84%) of 4-hydroxybenzothiophene
m.p. 76-78°C, content: 98%.

Example 2B

10 4-Hydroxybenzothiophene

[0053] An autoclave was charged under an argon flow with 27.34 g of 1-(2-thienyl)allyl acetate (0.150 mol, crude quality, see Example 1), 28.4 ml of acetic anhydride (30.6 g, 0.30 mol), 42.0 ml of triethylamine (30.7 g, 30 mol), 23.6 mg of palladium acetate (0.105 mmol) and 0.264 g of triphenylphosphine (1.00 mmol), all with aid of 53 ml of toluene.

15 Then the autoclave was sealed, evacuated twice under slow stirring (150 rpm) to 0.2 bar and pressurized with 8 bar of argon, then pressurized three times with 20 bar of carbon monoxide and vented, and finally pressurized with 50 bar of carbon monoxide. The reaction mixture was stirred (500 rpm) and heated at 120°C and the carbonylation carried out at 50 bar constant total pressure for 6 h. After cooling, the autoclave was vented and the CO atmosphere was exchanged by evacuating to ca. 0.2 bar and pressurizing 8 bar of argon four times. The resulting dark solution was poured into a

20 0.51 flask containing 120 ml of ice water and the biphasic solution was stirred for 1 h at room temperature. The aqueous phase was extracted in a separatory funnel with 80 ml of toluene and then combined organic phases were washed with 3 x 30 ml, a total of 90 ml of deionized water and reduced to a total weight of 46 g by rotary evaporation (50°C/60 mbar).

[0054] The residue containing the crude acetate was filtered through 17 g of silica gel ($\varnothing=3$ cm) and the filter washed with 150 ml of toluene. The combined organic phases were reduced to a total weight of 40 g by rotary evaporation and

25 transferred to a 0.351 glass flask under argon with aid of 20 ml of toluene. After addition of 82 ml of 4N sodium hydroxide (328 mmol) the mixture was stirred intensively (1200 rpm) at 50° for 1.5 h and then after cooling transferred in a 0.5 l separatory funnel. After removal of the organic layer, the dark aqueous phase was extracted with 80 ml of toluene and the combined organic phases were back-extracted with 2 x 20 ml, a total of 40 ml of deionized water. The combined aqueous phases were treated with 1.0 g of charcoal, stirred at room temperature for 5 min under argon and filtered

30 through a Speedex layer. The filter cake was rinsed three times with 20 ml, a total of 60 ml of deionized water. The clear brown, combined phases were concentrated until no more toluene distilled, then after cooling to 5°C in an ice bath 75 ml of 25% HCl were added under argon during 35 min, whereas the temperature was kept under 15° with aid of an ice bath. The resulting thick crystalline suspension was stirred for 1 h in an ice bath (internal temperature 2-3°C) and filtered on a sintered glass filter. The filter cake was washed three times with 50 ml, a total of 150 ml of ice cold water and dried

35 on the rotavapor at 50°C/l mbar to constant weight.
Yield: 16.5 g (73%) of 4-hydroxybenzothiophene as brown crystals-m.p. 75-76°C, content: 95%.

Example 3

40 Variation of Phosphorus Ligands

[0055] 4.93 mg of palladium acetate and 57.57 mg of triphenylphosphine in 10 ml of toluene were stirred for 1 h in a glove-box ($O_2 < 1$ ppm). A 35 ml autoclave was charged with 0.40 g of distilled 1-(2-thienyl)allyl acetate, 0.42 ml of acetanhydride, 0.62 ml of triethylamine and 1.0 ml of the catalyst solution described above. The autoclave was conditioned

45 with 30 bar of CO and pressurized with 70 bar of CO. The cyclocarbonylation was carried out at 120°C for 2 h. GC-analysis revealed a conversion of 96% with a content of 4-acetoxybenzothiophene of 91%.

A) Examples 3.1 - 3.6:

50 [0056] According to Example 3, table 1 summarizes the following experiments which were performed with phosphorus ligands other than triphenylphosphine.

Table 1:

Example	P-ligand ^{b)}	% conversion ^{a)}	% Content of 4-Acetoxybenzothiophene ^{a)}
3.1	PPh(3,5-tBu-Ph) ₂	92	87
3.2	P(3,5-tBu-Ph) ₃	93	88

Table continued

Example	P-ligand ^{b)}	% conversion ^{a)}	% Content of 4-Acetoxybenzothiophene ^{a)}
3.3	AMPHOS	99	95
3.4	NMDPP	98	94
3.5	P(2-Furyl) ₃	96	90
3.6	P(o-DMA-Ph) ₃	13	12
^{a)} Determined via GC (area-%). ^{b)} See structures in Scheme 1. ^{c)} P/Pd = 2			

B) Examples 3.7 - 3.23:

[0057] The following additional examples 3.7 to 3.23 were performed with further phosphorus ligands. The reaction were performed according to the description given above. However, the autoclave was pressurized with 50 bar CO and the cyclocarbonylation reaction was carried out at 90°C for 16 - 18 h.

Table 2:

Example No.	P-Ligand ^{b)}	P/Pd ^{c)}	% conversion ^{a)}	% Content ^{a)}
3.7	PPh(3,5-tBu-Ph) ₂	2	>99	94
3.8	P(3,5-tBu-Ph) ₃	2	>99	97
3.9	PAMP	2	>99	92
3.10	MOP	2	>99	92
3.11	P(2-Furyl) ₃	10	>99	85
3.12	TROPP-Ph	6	>99	95
3.13	PPh(Diphol)	6	>99	88
3.14	(S,S)-DDPPI	2	>99	92
3.15	DPEphos	2	99	40
3.16	DPPM	2	40	30
3.17	DIOP	4	99	72
3.18	P(O-nC ₄ H ₉) ₃	6	99	86
3.19	Diphol-DIOP	2	99	71
3.20	DPPF	2	>99	81
3.21	TPP-ox-Ph	1	85	44
3.22	P(m-Tol) ₃	6	>99	83
3.23	P(n-Bu) ₃	6	95	70
^{a)} % Content of 4-acetoxybenzothiophene, determined via GC (Area%). ^{b)} See structures in Scheme 1. ^{c)} Phosphorus-to-palladium molar ratio.				

Example 4

Cyclocarbonylation reactions: CO pressure and S/Pd ratio

[0058] 6.0 g of distilled 1-(2-thienyl)allyl acetate were reacted for 4 h as described in Example 2 A with 6.2 mg of palladium acetate, 72.1 mg of triphenylphosphine, 6.3 ml of acetic anhydride and 9.3 ml of triethylamine present. GC-analysis revealed a conversion of 98% with a content of 4-acetoxybenzothiophene of 94 %.

Examples 4.1 - 4.7:


[0059] According to Example 4 table 2 summarizes experiments performed under different reaction conditions (CO pressure and S/Pd ratio).

Table 2:

Example No.	Pco [bar] ^{a)}	S/Pd	T [°C]	% yield ^{b)}		
				2h	4h	6h
4.1	70	1200	120	99	-	-
4.2	70	"	100	90	98	n.d.
4.3	40	"	120	96	99	>99
4.4	20 ;	"	"	67	88	89
4.5	70	1500	"	97	>99	-
4.6	50	"	"	n.d.	n.d.	>99
4.7	40	"	"	92	97	98
a) determined at RT.						
b) Determined via GC (area-%).						

Example 5**4-[2-(Benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole**

[0060] 218 g (1.45 mol) of 4-hydroxy-benzothiophene and 511 g (1.82 mol) of mesylate of Formula V



were dissolved in 5.4 l of DMF, followed by addition of 555 g (4.02 mol) of potassium carbonate (dry). The reaction mixture was stirred at 100 to 105 °C for 6 to 8 hours. The resulting suspension was cooled to 5 °C and 7 l of water was added. The suspension was stirred at 5 °C for 30 minutes. The precipitate was filtered with suction and washed with 550 ml of DMF/water (1:1) and 1,1 l water. The precipitate was stirred at 0 to 5 °C in 1 l of MEK (methyl ethyl ketone) for 30 minutes. Then the precipitate was filtered with suction and dried at 50 °C.

Yield: 365 g (= 75 %) 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyloxazole.

m.p. 126 °C/129-131 °C

Example 6**5-Methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole**

[0061] 286 g (0.853 mol) of 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyloxazole were suspended in 6.3 l of glacial acetic acid. Temperature was raised to 60 °C. The resulting clear solution was cooled to 25 °C. 132 ml (3.18 mol) of 100 % nitric acid were added within 3 minutes. The reaction mixture was cooled below 30 °C. After crystallisation the suspension was stirred at 18 to 20 °C for 1 hour. The precipitate was filtered with suction and washed with 2 x 600 ml of tert-butyl methyl ether. The residue was suspended in 4 l of acetic ester for 15 minutes. 200 g (1.9 mol) of sodium carbonate in 3 l water were added. The resulting suspension was stirred for 1 hour. Acetic ester was distilled off followed by addition of 2 l of water. The suspension was stirred for 30 minutes. The precipitate was filtered with suction, washed with water and dried (50 °C, 24 hours).

Yield: 210 g of 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyloxazole (= 70 %).

m.p. 149 - 151 °C.

Example 7**5-Methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole**

- 5 **[0062]** 50 g (1.052 mol) of 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole were solved in 1 l of THF at 20 to 25 °C. 75 ml Lewatit M 600 (OH-form) (Bayer AG) were washed with about 100 ml THF, added to the 5-methyl-4-[2-(7-nitrobenzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole solution and stirred at room temperature for 1 hour. Then the Lewatit M 600 material was filtered with suction and washed with 100 ml THF. 12.5 g of Raney nickel were added to the combined THF solutions followed by hydrogenation of the 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole at standard pressure. The temperature of the reaction mixture should not exceed 35 to 40 °C. Hydrogen pressure was increased to 6 bar within 6 hours. After hydrogenation the reaction mixture was stirred for 1 hour. Then the catalyst was filtered with suction, the THF was distilled off, 180 ml ethanol was added and the residue was boiled out for 30 minutes. The reaction mixture was stirred at 0 °C for 1 hour. The precipitate was filtered with suction and the residue was washed with 25 ml ethanol and dried for 24 hours at 50°C (vacuum).
- 10
- 15 Yield: 42.4 g of 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyloxazole (= 92 %).
m.p. 122 -126 °C.

Example 8

- 20 **Methyl-2-bromo-3-[4-[2-(S-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate**

- [0063]** 320 g (0.91 mol) of 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole were solved in 6.41 acetone. Within 30 sec one third of a 320 ml (2.74 mol) of 48 % HBr in 900 ml water were added. After cooling to 0 to 4 °C and crystallization the suspension was stirred at 0 to 4 °C for 1 hour. Then the remaining 48 % HBr solution in water was added within 15 minutes at 0 to 4 °C and stirred at this temperature for 15 minutes followed by addition of 63.9 g (0.93 mol) sodium nitrite in 180 ml water within 15 minutes at 3 to 5 °C and stirring for 30 minutes at 3 to 5 °C. 1230 ml of methacrylate $\text{CH}=\text{CHCOOCH}_3$ (13.6 mol) was added to this reaction mixture at 10 to 14 °C followed by addition of 3.2 g Cu(I)bromide. Temperature was increased to 20 to 25 °C within 30 minutes followed by stirring at this temperature for 1 hour and 10 minutes at 30 °C. 1.81 of water was added to the reaction mixture followed by distillation of acetone/methacrylate $\text{CH}=\text{CHCOOCH}_3$ at a temperature of 40 °C. The final volume was about 21.1 l of water was added to separate the remaining methacrylate $\text{CH}=\text{CHCOOCH}_3$. The final volume was about 21. The black precipitate was solved by addition of 4.51 of acetic ester and stirring for 15 minutes. The two phase reaction mixture was filtered and the aqueous phase was extracted with 2 l of acetic ester. After extraction with 2 l of an aqueous 2 % NaCl-solution the acetic ester solutions were combined and distilled. 2 l of acetic ester was added to the residue and again distilled. 3 l of ethanol were added to the residue and boiled. 15 g of activated charcoal were added and stirred for 15 minutes. After filtration and cooling to room a precipitate was formed. The suspension was stirred for 1 hour at room temperature and an additional hour at 0 °C. After washing with cold ethanol the precipitate was dried for 24 hours at 50°C (vacuum). Yield: 310 g of methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate (= 68 %).
- 30
- 35
- 40 m.p. 97 to 99 °C.

Example 9

- 45 **2-Imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methyl-thiazolidine-4-one**

- [0064]** 190 g (0.380 mol) of 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole were suspended in 2.85 l of ethanol. 31.6 g (0.415 mol) of thiourea and 34.8 g of sodium acetate were added. After boiling for about 18 hours (reflux) the reaction mixture was cooled to 0 to 4 °C and stirred for 1.5 hours at this temperature. The precipitate was filtered with suction and washed twice with 250 ml cold ethanol. 1.9 l of water was added to the residue, the mixture was stirred for 10 minutes and the precipitate was filtered with suction and dried for 24 hours at 80 °C (vacuum). Yield: 147 g of 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methyl-thiazolidine-4-one (84 %); m.p: 224 - 227 °C.
- 50
- 55

Example 10

5-[4-[2-(5-Methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione

[0065] 283.3 g (0.61 mol) of 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methyl-thiazolidine-4-one were suspended in 2.83 l ethanol. 2.83 l of 2 N hydrochloric acid were added. The resulting suspension was stirred for 18 hours (reflux). The suspension was cooled for 1 hour to 0 to 4 °C and was stirred for another 2 hours at this temperature. The precipitate was filtered with suction and washed twice with 285 ml of ethanol. 2.83 l of water was added to the residue, the suspension was stirred for 30 minutes, the precipitate was filtered with suction and washed with 21 of water. The precipitate was dried for 24 hours at 80 °C and then solved in 545 ml DMF (at 85 to 90 °C). 4.95 l of ethanol (25 °C) were added to the solution. The resulting suspension was stirred for 2 hours at 0 to 4 °C. The precipitate was filtered with suction, washed with 270 ml of cold ethanol and dried at 80 °C for 24 hours (vacuum).

Yield: 246 g of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione (87%);
MP: 224 - 227 °C.

Example 11


5-[4-[2-(5-Methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt

[0066] (5-[4-[2-(5-Methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzo[b]thiophen-7-ylmethyl]-2,4-thiazolidinedione) (5.8 g) was dissolved in hot THF (87 ml). A solution of sodium hydroxide (0.5 g) in water (6 ml) was added, and the solution was cooled to room temperature. Another portion (87 ml) of THF was given to the solution, and after a short time a crystallization was observed. 150 ml of the solvent was distilled off in the heat. The suspension was cooled to about 0°C and was stirred for further 2 hours. The solid was filtered and dried at 80°C.

Yield: 5.6 g of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt (93%);
MP: > 300°C (decomposition).


Claims

1. A process for the preparation of compounds of Formula I



I

comprising cyclocarbonylation of a compound of Formula II




II

wherein -OR is a group of formulae -O-(CO)-R', -O-(CO)-O-R'' or -O-(PO)-(OR'')₂, wherein R' is alkyl, perfluoro-C₁₋₂₀-alkyl, aryl, R'' is alkyl, aryl or benzyl or the group -OR is halogen or an aryloxy group;
followed by saponification.


2. The process according to claim 1, wherein the cyclocarbonylation reaction is carried out in the presence of a base and a catalyst comprising a transition metal compound and a ligand.


3. The process according to any of claims 1- 2, wherein the transition metal compound is a palladium salt.
4. The process according to claim 3, wherein the transition metal compound is selected from a group consisting of Pd(OAc)₂, Pd₂dba₃, PdCl₂, Pd₂Cl₂(π -allyl)₂, PdCl₂(NCMe)₂, [Pd(NCMe)₄](BF₄)₂ or Pd/C.
5. The process according to claim 4, wherein the palladium compound is Pd(OAc)₂.
6. The process according to claims 1- 5, wherein the ligand is P(R¹)(R²)(R³) or (R¹)(R²)P-(X)-P(R¹)(R²) wherein R¹, R² and R³ each independently are C₁₋₈-alkyl, cyclohexyl, benzyl, naphthyl, 2- or 3-pyrrolyl, 2- or 3-furyl, 2- or 3-thiophenyl, 2- or 3- or 4-pyridyl, phenyl or phenyl which is substituted by C₁₋₄-alkyl, C₁₋₄-alkoxy, halogen, trifluoromethyl, lower alkyldenedioxy or phenyl and X is binaphthyl, 6,6'-dimethyl- or 6,6'-dimethoxybiphenyl-2,2'-diyl, or one of the groups -(CH₂)_n-, -CH₂CH₂-P(C₆H₅)-CH₂CH₂-.




and n is a number of 1- 8.

7. The process according to claims 1- 5, wherein the ligand is selected from a group consisting of triphenylphosphine, and






and



8. The process according to claim 1 - 7, wherein the ligand is triphenylphosphine,




9. The process according to claims 1 - 8, wherein the cyclocarbonylation reaction is carried out in the presence of a

base selected from the group consisting of tri-alkyl-amines, di-alkyl-aryl-amines, pyridines, alkyl-N-piperidines, sodium hydroxide, potassium hydroxide or salts of carbonic acids.

10. The process according to claims 1- 9, wherein the cyclocarbonylation reaction is carried out in the presence of triethylamine.

11. The process according to claims 1 - 10, wherein the compounds of Formula II are prepared by reaction of the thiophene carbaldehyde of Formula III



III

with a reagent of the formula vinyl-metal-X with -metal-X being -MgCl, -MgBr, -MgI or -Li, followed by reaction with an acid derivative selected from a group consisting of (R'-CO)₂O, R''O-(CO)-Cl, Cl-(PO)(OR'')₂ or R'''-(CO)-Hal, wherein R' is alkyl, perfluoro-C₁₋₂₀-alkyl, aryl, R'' is alkyl, aryl or benzyl and Hal is Cl or Br.

12. The process according to claim 11, wherein the acid derivative is acetanhydrid.


13. The process according to claims 1 - 12, wherein the vinyl-metal-X-reagent is vinylmagnesium chloride.

14. The process according to claims 1 - 13, wherein the compound of Formula II is prepared by reaction of the thiophene carbaldehyde by reaction of vinylmagnesium chloride followed by reaction with acetanhydride.

15. The process according to claims 1 - 14, wherein the compound of Formula I is 4-hydroxybenzothiophene.

16. The process according to claims 1 - 15, wherein the saponification reaction is carried out in a biphasic mixture of sodium hydroxide in toluene or in a homogeneous mixture of sodium methylate in methanol.

17. The process according to claims 1 - 16, wherein the compound of Formula I is converted into 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole by reaction of a compound of Formula I with a mesylate of Formula V



under basic conditions.

18. The process according to claim 17, wherein 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole is converted into 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole by nitration.

19. The process according to claim 18, wherein 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole is converted into 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole by hydrogenation.

20. The process according to claim 19, wherein 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole is converted into methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate by reaction with HHal/NaNO₂ followed by reaction with CH=CHCOOCH₃/Cu(I)Hal, wherein Hal is Br or Cl.

21. The process according to claim 20, wherein methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate is converted into 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothi-


ophene-7-yl]-methyl-thiazolidine-4-one by reaction with thiourea.

22. The process according to claim 21, wherein 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methyl-thiazolidine-4-one is converted into 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione by reaction under acid conditions.

23. The process according to claim 22, wherein 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione is converted into 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt by reaction under basic conditions.


24. The process according to claims 1 to 23 for the preparation of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione comprising

a) conversion of a compound of Formula I into 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole by reaction of a compound of Formula I



I

with a mesylate of Formula V



V

under basic conditions; followed by

b) nitration of 4-[2-(benzothiophene-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazole to give 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole;

c) hydrogenation of 5-methyl-4-[2-(7-nitro-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole to give 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole; followed by

d) reaction of 5-methyl-4-[2-(7-amino-benzothiophene-4-yloxy)-ethyl]-2-phenyl-oxazole with $\text{HHal}/\text{NaNO}_2$ and $\text{CH}=\text{CHCOOCH}_3/\text{Cu(I)Hal}$, wherein Hal is Br or Cl to give methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate; followed by

e) reaction of methyl-2-bromo-3-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-propionate with thiourea to give 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methylthiazolidine-4-one; followed by

f) reaction of 2-imino-5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-yl]-methylthiazolidine-4-one under acid conditions to give 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione;

g) and optionally followed by reaction of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione under basic conditions to give 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt.


25. The use of a process according to any of claims 1- 24 for the preparation of 5-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]-7-benzothiophenylmethyl]-2,4-thiazolidinedione.

26. The use of a process according to any of claims 1 - 24 for the preparation of 5-[4-[2-(5-methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt.

27. 5-[4-[2-(5-Methyl-2-phenyl-oxazole-4-yl)-ethoxy]-benzothiophene-7-methyl]-2,4-thiazolidinedione-Na-salt.


Patentansprüche

1. Verfahren zur Herstellung von Verbindungen der Formel I



I

umfassend die Cyclocarbonylierung einer Verbindung der Formel II



II

worin -OR eine Gruppe der Formeln $-O-(CO)-R'$, $-O-(CO)-O-R''$ oder $-O-(PO)-(OR''')_2$ ist, worin R' Alkyl, Perfluor- C_{1-20} -alkyl, Aryl ist, R'' Alkyl, Aryl oder Benzyl ist oder die Gruppe -OR Halogen oder eine Aryloxygruppe ist; gefolgt von Verseifung.


2. Verfahren nach Anspruch 1, wobei die Cyclocarbonylierungsreaktion in Gegenwart einer Base und eines Katalysators, umfassend eine Übergangsmetallverbindung und einen Liganden, durchgeführt wird.

3. Verfahren nach einem der Ansprüche 1 bis 2, wobei die Übergangsmetallverbindung ein Palladiumsalz ist.


4. Verfahren nach Anspruch 3, wobei die Übergangsmetallverbindung aus der Gruppe ausgewählt ist, bestehend aus $Pd(OAc)_2$, Pd_2dba_3 , $PdCl_2$, $Pd_2Cl_2(\pi\text{-allyl})_2$, $PdCl_2(NCMe)_2$, $[Pd(NCMe)_4](BF_4)_2$ oder Pd/C .

5. Verfahren nach Anspruch 4, wobei die Palladiumverbindung $Pd(OAc)_2$ ist.

6. Verfahren nach den Ansprüchen 1 bis 5, wobei der Ligand $P(R^1)(R^2)(R^3)$ oder $(R^1)(R^2)P(X)P(R^1)(R^2)$ ist, worin R^1 , R^2 und R^3 jeweils unabhängig voneinander C_{1-8} -Alkyl, Cyclohexyl, Benzyl, Naphthyl, 2- oder 3-Pyrrolyl, 2- oder 3-Furyl, 2- oder 3-Thiophenyl, 2- oder 3- oder 4-Pyridyl, Phenyl oder Phenyl, das mit C_{1-4} -Alkyl, C_{1-4} -Alkoxy, Halogen, Trifluormethyl, Niederalkyldendioxy oder Phenyl substituiert ist, sind und X Binaphthyl, 6,6'-Dimethyl- oder 6,6'-Dimethoxybiphenyl-2,2'-diyl oder eine der Gruppen $-(CH_2)_n$, $-CH_2CH_2-P(C_6H_5)-CH_2CH_2-$,



oder




IV

ist und n eine Zahl von 1 bis 8 ist.


7. Verfahren nach den Ansprüchen 1 bis 5, wobei der Ligand aus der Gruppe ausgewählt ist, bestehend aus Triphenylphosphin und




P(o-DMA-Ph)₃




PPh(3,5-tBu-Ph)₂




P(3,5-tBu-Ph)₃




P(2-Furyl)₃




NMDPP




AMPHOS



DIOP




PAMP




MOP


und




(S,S)-DDPPI




DPEphos




DPPM




TPP-ox-Ph




P(m-Tol)₃




DPPF



TROP-Ph




PPh(Diphol)




Diphol-DIOP

8. Verfahren nach Anspruch 1 bis 7, wobei der Ligand Triphenylphosphin,

PPh(3,5-tBu-Ph)₂

oder


P(3,5-tBu-Ph)₃

ist.

9. Verfahren nach den Ansprüchen 1 bis 8, wobei die Cyclocarbonylierungsreaktion in Gegenwart einer Base durchgeführt wird, ausgewählt aus der Gruppe, bestehend aus Tri-alkylaminen, Di-alkyl-aryl-aminen, Pyridinen, Alkyl-N-piperidinen, Natriumhydroxid, Kaliumhydroxid oder Salzen von Kohlensäuren.

10. Verfahren nach den Ansprüchen 1 bis 9, wobei die Cyclocarbonylierungsreaktion in Gegenwart von Triethylamin durchgeführt wird.

11. Verfahren nach den Ansprüchen 1 bis 10, wobei die Verbindungen der Formel II durch die Umsetzung des Thiophencarbaldehyds der Formel III



III

mit einem Reagens der Formel Vinyl-Metall-X hergestellt werden, wobei -Metall-X -MgCl-, -MgBr, -MgI oder -Li ist, gefolgt von der Umsetzung mit einem Säurederivat, ausgewählt aus der Gruppe, bestehend aus (R'-CO)₂O, R''O-(CO)-Cl, Cl-(PO)(OR'')₂ oder R''-(CO)-Hal, worin R' Alkyl, Perfluor-C₁₋₂₀-alkyl, Aryl ist, R'' Alkyl, Aryl oder Benzyl ist und Hal Cl oder Br ist.

12. Verfahren nach Anspruch 11, wobei das Säurederivat Acetanhydrid ist.


13. Verfahren nach den Ansprüchen 1 bis 12, wobei das Vinyl-Metall-X-Reagens Vinylmagnesiumchlorid ist.

14. Verfahren nach den Ansprüchen 1 bis 13, wobei die Verbindung der Formel II durch die Umsetzung des Thiophencarbaldehyds durch die Umsetzung von Vinylmagnesiumchlorid, gefolgt von der Umsetzung mit Acetanhydrid hergestellt wird.

15. Verfahren nach den Ansprüchen 1 bis 14, wobei die Verbindung der Formel I 4-Hydroxybenzothiopen ist.

16. Verfahren nach den Ansprüchen 1 bis 15, wobei die Verseifungsreaktion in einem zweiphasigen Gemisch aus Natriumhydroxid in Toluol oder in einem homogenen Gemisch aus Natriummethylat in Methanol durchgeführt wird.

17. Verfahren nach den Ansprüchen 1 bis 16, wobei die Verbindung der Formel I zu 4-[2-(Benzothiophen-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazol durch die Umsetzung einer Verbindung der Formel I mit einem Mesylat der Formel V



unter basischen Bedingungen umgewandelt wird.

18. Verfahren nach Anspruch 17, wobei 4-[2-(Benzothiophen-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazol zu 5-Methyl-4-[2-(7-nitro-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol durch Nitrierung umgewandelt wird.

19. Verfahren nach Anspruch 18, wobei 5-Methyl-4-[2-(7-nitro-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol zu 5-Methyl-4-[2-(7-amino-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol durch Hydrierung umgewandelt wird.

20. Verfahren nach Anspruch 19, wobei 5-Methyl-4-[2-(7-amino-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol zu Methyl-2-brom-3-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-propionat durch die Umsetzung mit $\text{HHal}/\text{NaNO}_2$, gefolgt von der Umsetzung mit $\text{CH}=\text{CHCOOCH}_3/\text{Cu(I)Hal}$, worin Hal Br oder Cl ist, umgewandelt wird.


21. Verfahren nach Anspruch 20, wobei Methyl-2-brom-3-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-propionat zu 2-Imino-5-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-methyl-thiazolidin-4-on durch die Umsetzung mit Thioharnstoff umgewandelt wird.

22. Verfahren nach Anspruch 21, wobei 2-Imino-5-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-methyl-thiazolidin-4-on zu 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion durch die Umsetzung unter sauren Bedingungen umgewandelt wird.


23. Verfahren nach Anspruch 22, wobei 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion zu 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion-Na-Salz durch die Umsetzung unter basischen Bedingungen umgewandelt wird.

24. Verfahren nach den Ansprüchen 1 bis 23 zur Herstellung von 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion, umfassend

a) Umwandlung einer Verbindung der Formel I in 4-[2-(Benzothiophen-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazol durch die Umsetzung einer Verbindung der Formel I



mit einem Mesylat der Formel V



5

10

15

20

25

- unter basischen Bedingungen; gefolgt von
- b) Nitrierung von 4-[2-(Benzothiophen-4-yloxy)-ethyl]-5-methyl-2-phenyl-oxazol, um 5-Methyl-4-[2-(7-nitro-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol zu erhalten;
- c) Hydrierung von 5-Methyl-4-[2-(7-nitro-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol, um 5-Methyl-4-[2-(7-amino-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol zu erhalten; gefolgt von
- d) Umsetzung von 5-Methyl-4-[2-(7-amino-benzothiophen-4-yloxy)-ethyl]-2-phenyl-oxazol mit HHal/NaNO_2 und $\text{CH=CHCOOCH}_3/\text{Cu(I)Hal}$, worin Hal Br oder Cl ist, um Methyl-2-brom-3-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-propionat zu erhalten; gefolgt von
- e) Umsetzung von Methyl-2-brom-3-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-propionat mit Thioharnstoff, um 2-Imino-5-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-methyl-thiazolidin-4-on zu erhalten; gefolgt von
- f) Umsetzung von 2-Imino-5-[4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-yl]-methyl-thiazolidin-4-on unter sauren Bedingungen, um 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion zu erhalten;
- g) und gegebenenfalls gefolgt von der Umsetzung von 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidinon unter basischen Bedingungen, um 5-[4-[2-(S-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion-Na-Salz zu erhalten.

25. Verwendung eines Verfahrens nach einem der Ansprüche 1 bis 24 zur Herstellung von 5-[4-[2-(5-Methyl-2-phenyl-4-ethoxy)-benzothiophen-7-methyl]-2,4-thiazolidindion.

30

26. Verwendung eines Verfahrens nach einem der Ansprüche 1 bis 24 zur Herstellung von 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion-Na-Salz.


27. 5-[4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzothiophen-7-methyl]-2,4-thiazolidindion-Na-Salz.

35

Revendications

1. Procédé de préparation de composés de formule I

40




45

I

comprenant la cyclocarbonylation d'un composé de formule II

50




55

II



dans laquelle -OR est un groupe de formule -O-(CO)-R', -O-(CO)-O-R'' ou -O-(PO)-(OR'')₂ où R' est un groupe alkyle, perfluoralkyle en C₁ à C₂₀, aryle, R'' est un groupe alkyle, aryle ou benzyle ou le groupe -OR est un halogène ou un groupe aryloxy ;
suivie d'une saponification.




2. Procédé selon la revendication 1, dans lequel on effectue la réaction de cyclocarbonylation en présence d'une base et d'un catalyseur comprenant un composé d'un métal de transition et un ligand.
3. Procédé selon l'une quelconque des revendications 1 et 2, dans lequel le composé de métal de transition est un sel de palladium.
4. Procédé selon la revendication 3, dans lequel on choisit le composé de métal de transition dans un groupe constitué de Pd(OAc)₂, Pd₂dba₃, PdCl₂, Pd₂Cl₂(π-allyl)₂, PdCl₂(NCMe)₂, [Pd(NCMe)₄](BF₄)₂ ou Pd/C.
5. Procédé selon la revendication 4, dans lequel le composé de palladium est Pd(OAc)₂.
6. Procédé selon les revendications 1 à 5, dans lequel le ligand est P(R¹)(R²)(R³) ou (R¹)(R²)P-(X)-P(R¹)(R²) où R¹, R² et R³ sont chacun indépendamment un groupe alkyle en C₁ à C₈, cyclohexyle, benzyle, naphthyle, 2- ou 3-pyrrolyle, 2- ou 3-furyle, 2- ou 3-thiophényle, 2- ou 3- ou 4-pyridyle, phényle ou phényle qui est substitué par un groupe alkyle en C₁ à C₄, alkoxy en C₁ à C₄, halogéno, trifluorométhyle, alkyldènedioxy inférieur ou phényle, et X est un groupe binaphtyle, 6,6'-diméthyl- ou 6,6'-diméthoxybiphényl-2,2'-diyle ou l'un des groupes -(CH₂)_n-, -CH₂CH₂-P(C₆H₅)-CH₂CH₂-.











et n est un nombre de 1 à 8.

7. Procédé selon les revendications 1 à 5, dans lequel on choisit le ligand dans le groupe constitué de triphénylphosphine et







et



8. Procédé selon les revendications 1 à 7, dans lequel le ligand est de la triphénylphosphine,




9. Procédé selon les revendications 1 à 8, dans lequel on effectue la réaction de cyclocarbonylation en présence d'une base choisie dans le groupe constitué de trialkylamines, dialkylarylamines, pyridines, alkyl-N-pipéridines, hydroxyde de sodium, hydroxyde de potassium ou sels d'acides carboniques.
10. Procédé selon les revendications 1 à 9, dans lequel on effectue la réaction de cyclocarbonylation en présence de triéthylamine.
11. Procédé selon les revendications 1 à 10, dans lequel on prépare les composés de formule II par réaction du thiophénecarbaldéhyde de formule III



III

avec un réactif de formule vinyl-métal-X, -métal-X représentant -MgCl, -MgBr, -MgI ou -Li, suivie d'une réaction avec un dérivé d'acide choisi dans le groupe constitué de $(\text{R}'\text{-CO})_2\text{O}$, $\text{R}''\text{O}(\text{CO})\text{-Cl}$, $\text{Cl}(\text{PO})(\text{OR}'')_2$ ou $\text{R}''\text{-(CO)-Hal}$ où R' est un groupe alkyle, perfluoralkyle en C_1 à C_{20} , aryle, R'' est un groupe alkyle, aryle ou benzyle et Hal est Cl ou Br.

12. Procédé selon la revendication 11, dans lequel le dérivé d'acide est de l'anhydride acétique.
13. Procédé selon les revendications 1 à 12, dans lequel le réactif vinyl-métal-X est du chlorure de vinylmagnésium.
14. Procédé selon les revendications 1 à 13, dans lequel on prépare le composé de formule II par réaction du thiophénecarbaldéhyde par réaction du chlorure de vinylmagnésium, suivie d'une réaction avec de l'anhydride acétique.
15. Procédé selon les revendications 1 à 14, dans lequel le composé de formule I est du 4-hydroxybenzothiophène.
16. Procédé selon les revendications 1 à 15, dans lequel on effectue la réaction de saponification dans un mélange biphasique d'hydroxyde de sodium dans du toluène ou dans un mélange homogène de méthylate de sodium dans du méthanol.
17. Procédé selon les revendications 1 à 16, dans lequel le composé de formule I est transformé en 4-[2-(benzothiophène-4-yloxy)-éthyl]-5-méthyl-2-phényl-oxazole par réaction d'un composé de formule I avec un mésylate de formule V



V

dans des conditions basiques.

18. Procédé selon la revendication 17, dans lequel le 4-[2-(benzothiophène-4-yloxy)-éthyl]-5-méthyl-2-phényl-oxazole est transformé en 5-méthyl-4-[2-(7-nitro-benzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole par nitration.

19. Procédé selon la revendication 18, dans lequel le 5-méthyl-4-[2-(7-nitro-benzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole est transformé en 5-méthyl-4-[2-(7-aminobenzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole par hydrogénéation.

20. Procédé selon la revendication 19, dans lequel le 5-méthyl-4-[2-(7-amino-benzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole est transformé en propionate de méthyl-2-bromo-3-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yle] par réaction avec HHal/NaNO_2 suivie d'une réaction avec $\text{CH=CHCOOCH}_3/\text{Cu(I)Hal}$ où Hal est Br ou Cl.


21. Procédé selon la revendication 20, dans lequel le propionate de méthyl-2-bromo-3-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yle] est transformé en 2-imino-5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yl]-méthyl-thiazolidine-4-one par réaction avec de la thio-urée.

22. Procédé selon la revendication 21, dans lequel la 2-imino-5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yl]-méthyl-thiazolidine-4-one est transformée en 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione par réaction dans des conditions acides.

23. Procédé selon la revendication 22, dans lequel la 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione est transformée en sel de sodium de 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione par réaction dans des conditions basiques.


24. Procédé selon les revendications 1 à 23 pour la préparation de 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione, comprenant

a) la transformation d'un composé de formule I en 4-[2-(benzothiophène-4-yloxy)-éthyl]-5-méthyl-2-phényl-oxazole par réaction d'un composé de formule V



I

avec un mésylate de formule V



V

dans des conditions basiques ; suivie de

b) la nitration du 4-[2-(benzothiophène-9-yloxy)-éthyl]-5-méthyl-2-phényl-oxazole pour obtenir du 5-méthyl-4-[2-(7-nitro-benzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole ;

c) l'hydrogénation du 5-méthyl-4-[2-(7-nitrobenzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole pour obtenir du 5-méthyl-4-[2-(7-aminobenzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole ; suivie de

d) une réaction du 5-méthyl-4-[2-(7-amino-benzothiophène-4-yloxy)-éthyl]-2-phényl-oxazole avec HHal/NaNO_2 et $\text{CH=CHCOOCH}_3/\text{Cu(I)Hal}$, où Hal est Br ou Cl, pour obtenir du propionate de méthyl-2-bromo-3-[4-[2-(5-mé-

thyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yle] ; suivie de
 e) une réaction du propionate de méthyl-2-bromo-3-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yle] avec de la thio-urée pour obtenir de la 2-imino-5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yl]-méthyl-thiazolidine-4-one ; suivie de

f) une réaction de la 2-imino-5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-yl]-méthyl-thiazolidine-4-one dans des conditions acides pour obtenir de la 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione ;

g) et éventuellement suivie d'une réaction de la 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione dans des conditions basiques pour obtenir un sel de sodium de 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione.

25. Utilisation d'un procédé selon l'une quelconque des revendications 1 à 24 pour la préparation de 5-[4-[2-(5-méthyl-2-phényl-4-oxazolyl)éthoxy]-7-benzothiophényméthyl]-2,4-thiazolidinedione.

26. Utilisation d'un procédé selon l'une quelconque des revendications 1 à 24 pour la préparation d'un sel de sodium de 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione.

27. Sel de sodium de 5-[4-[2-(5-méthyl-2-phényl-oxazole-4-yl)-éthoxy]-benzothiophène-7-méthyl]-2,4-thiazolidinedione.